

NATURAL RESOURCES CONSERVATION SERVICE CONSERVATION PRACTICE STANDARD

RIPARIAN FOREST BUFFER

(Acre)

CODE 391

DEFINITION

An area of predominantly trees and/or shrubs located adjacent to and up-gradient from watercourses or water bodies.

PURPOSES

- Create shade to lower **or moderate** water temperatures to improve habitat for aquatic organisms.
- Provide a source of detritus and large woody debris for aquatic and terrestrial organisms.
- Create wildlife habitat and establish wildlife corridors.
- Reduce excess amounts of sediment, organic material, nutrients and pesticides in surface runoff and reduce excess nutrients and other chemicals in shallow ground water flow.
- Provide a harvestable crop of timber, fiber, forage, fruit, or other crops consistent with other intended purposes.
- Provide protection against scour erosion within the floodplain.
- Restore natural riparian plant communities.
- Moderate winter temperatures to reduce freezing of aquatic over-wintering habitats.
- To increase carbon storage.

CONDITIONS WHERE PRACTICE APPLIES

On **stable** areas adjacent to permanent or intermittent streams, lakes, ponds, wetlands and areas with ground water recharge that are capable of supporting woody vegetation.

For areas with unstable banks refer to Streambank and Shoreline Protection - 580.

The riparian forest buffer is a component of a planned land management system including nutrient, pesticide, runoff, sediment, and erosion control practices.

CRITERIA

General Criteria Applicable To All Purposes

The location, layout, **width**, and density of the riparian forest buffer will **be selected to** accomplish the intended purpose and function.

All buffers will consist of a Zone 1 that begins at the normal water line, or at the top of the bank, and extends a minimum distance of 15 feet, measured horizontally on a line perpendicular to the water body.

Dominant **climax** vegetation will consist of existing, naturally regenerated, or planted trees and shrubs suited to the site and the intended purpose. ***Selection of native species will be a priority when feasible. Plantings will consist of two or more species with individual plants suited to the***

<p>Conservation practice standards are reviewed periodically, and updated if needed. To obtain the current version of this standard, contact the Natural Resources Conservation Service.</p>
--

seasonal variation of soil moisture status of individual planting sites. See figure 1. Plant types and species shall be selected based on their compatibility in growth rates, flooding tolerance, ability to produce large debris, and shade and wildlife values. Select species from Tables 1 and 2, Trees and Shrubs Suitable for Establishment located in Appendix I of this standard.

Other species not listed in Tables 1 and 2 may also be suitable. Consult WV Division of Forestry personnel, NRCS staff forester and/or biologist concerning the suitability of other species.




Figure 1. Plant adaptation to soil moisture.

Occasional removal of some tree and shrub products such as high value trees is permitted in Zone 1 provided the intended purpose is not compromised by the loss of vegetation or harvesting disturbance.

Harvesting operations in riparian forest buffers must adhere to current state and local regulations. Best Management Practices are outlined in Best Management Practices for Controlling Soil Erosion and Sedimentation from Logging Operations in West Virginia (WVDOF-TR-96-3).

Necessary site preparation and planting shall be done at a time and manner to insure survival and growth of selected species.

**NRCS, NHCP
August 2000**

Site preparation shall be sufficient for establishment and growth of selected species and done in a manner that does not compromise the intended purpose. **See Forest Site Preparation – 490.**

Only viable, high quality and adapted planting stock will be used.

For optimal carbon storage, select plant species that are adapted to the site to assure strong health and vigor and plant the full stocking rate for the site.

See Tree/Shrub Establishment – 612 for care, handling, and planting requirements of planting stock.

The method of planting for new buffers shall include hand or machine planting techniques and be suited to achieving proper depths and placement of planting stock roots.

An adequate nearby or adjacent seed source must be present when using natural regeneration to establish a buffer.

Livestock shall be controlled or excluded as necessary to achieve and maintain the intended purpose. **See Use Exclusion - 472.**

Fenced water course crossings and livestock watering areas shall be located and sized to minimize impact to the riparian buffer vegetation and function.

Harmful pests present on the site will be controlled or eliminated as necessary to achieve and maintain the intended purpose. **See Pest Management -595.**

Comply with applicable federal, state and local laws and regulations during the installation, operation (including harvesting activities) and maintenance of this practice.

Additional Criteria To Reduce Excess Amounts of Sediment, Organic Material, Nutrients and Pesticides in Surface Runoff and Reduce Excess Nutrients and Other Chemicals in Shallow Ground Water Flow

An additional strip or area of land, Zone 2, will begin at the edge and up-gradient of Zone 1

**NRCS, WV
February 2003**

and extend a minimum distance of 20 feet, measured horizontally on a line perpendicular to the water body. The minimum combined width of Zones 1 and 2 will be 100 feet or 30 percent of the flood plain whichever is less, but not less than 35 feet.

A flood plain is defined as the area adjacent to a river or stream that is built of alluvial sediments that are associated with present depositional activity. Figure 2 illustrates examples of Zone 1 and Zone 2 widths for watercourses and water bodies.

Criteria for Zone 1 shall apply to Zone 2 except that removal of tree and shrub products such as timber, nuts, fruit and forbs is permitted and encouraged on a periodic and regular basis provided the intended purpose is not compromised by loss of vegetation or harvesting disturbance.

Zone 2 will be expanded in high nutrient, sediment, and animal waste application areas, where the contributing area is not adequately treated or where an additional level of protection is desired.

Concentrated flow erosion shall be controlled in the up-gradient area immediately adjacent to Zone 2 prior to establishment of the riparian forest buffer. See Figure 3.




Figure 3. Control of concentrated flow erosion.

A Zone 3 shall be added to the riparian buffer when adjacent to cropland or other sparsely vegetated or highly erosive areas to filter sediment, address concentrated flow erosion, and maintain sheet flow. The Filter Strip - 393 standard shall be used to design Zone 3.

For areas other than those described above, or where the primary purpose of the riparian forest buffer is wildlife habitat, refer to Riparian Herbaceous Cover - 390 for more information concerning Zone 3.

Additional Criteria To Create Shade To Lower Or Moderate Water Temperatures

A buffer for lowering warm-season water temperatures shall consist of at least Zone 1.

Buffers shall be maintained or established on the south and west sides of water bodies insofar as practical. The buffer canopy shall be established to achieve at least 50 percent crown cover with the average canopy heights equal to or greater than the width of the watercourse or 30 feet for water bodies. See Figure 4.




Figure 4. Canopy height for water temperature control.

Buffer species shall include native riparian species of sufficient height and crown density potential. Place drooping or wide-crowned trees and shrubs nearest the watercourse or body. Shoreline or channel relief (e.g. deeply incised channels) and topographic shading should be taken into consideration when selecting species.




Figure 2. Examples of riparian forest buffer widths for watercourses and water bodies.

Additional Criteria To Provide A Source of Detritus And Large Woody Debris

Within Zone 1, as a minimum, establish, favor, or manage species capable of producing stems and limbs of sufficient size to provide an eventual source of large woody debris (> 10 inches in diameter) for instream habitat for fish and aquatic organisms.

Additional Criteria To Provide Habitat For Aquatic Organisms And Terrestrial Wildlife

Width of Zone 1 and/or Zone 2 will be expanded to meet the minimum requirements of the wildlife or aquatic species and associated communities of concern.

Establish plant communities that address the target wildlife needs and existing resources in the watershed. ***See Wildlife Wetland Habitat Management -644 and Wildlife Upland Habitat Management - 645.***

CONSIDERATIONS

Wide widths (75 feet or more) are preferred. They are more effective for the listed purposes and more feasible to manage. Narrower widths within this standard recognize the value of streamside land for farming and limited bottomland acreage in many locations.

Complex ownership patterns of riparian areas may require group planning for proper buffer design, function, and management.

The widths below are a guide for selected wildlife species. They include the sum of buffer widths on one or both sides of water courses or water bodies and may extend beyond riparian boundaries.

Species:	Desired Width in feet:
Bald eagle, wood ducks, heron rookery	600
Turkey, pileated woodpecker	450
Beaver, black duck, grouse, trout	300
Deer	200
Song birds, mallard	165
Bass, woodcock	100

The severity of bank erosion and its influence on existing or potential riparian trees and shrubs should be assessed. Watershed-level treatment or bank stability activities may be needed before establishing a riparian forest buffer.

When concentrated flow erosion and sedimentation cannot be controlled vegetatively, consider structural or mechanical treatments.

Where animal waste is a problem, management systems must be applied to the adjacent field. Forest buffers should not be planned as substitute practices.

Use of this practice without other nutrient, pesticide, sediment, and erosion control practices can result in adverse impacts on buffer vegetation and stream hydrology. The expected adverse impacts could be high maintenance costs, need for re-establishment of vegetation, and delivery of excess nutrients, sediment, and other potential pollutants through the buffer by concentrated flows.

Joining new buffers with existing buffers increases the continuity of cover and will further moderate water temperatures. A mix of species with growth forms that are

tall and wide-crowned or drooping will increase moderation effects.

For watercourses, buffers established on both sides will enhance riparian values.

Favor tree and shrub species that are non-invasive, or have multiple values such as those suited for timber, biomass, nuts, fruit, browse, nesting, aesthetics and tolerance to locally used herbicides. ***Consider species that resprout when establishing the vegetation nearest the watercourses or bodies. For detritus and large woody debris, use species that meet the specific requirements of fish and other aquatic organisms for food, habitat, migration and spawning.***

Use recommendations from regional or other large-scale evaluations and plans when designing, locating, and connecting buffers for indicator and/or target species of wildlife, fish, and other aquatic organisms.

Tree and shrub species, which may be alternate hosts to undesirable pests, should be avoided. Species diversity should be considered to avoid loss of function due to species-specific pests.

The location, layout and density of the buffer should compliment natural features, and mimic natural riparian forests.

Consider the type of human use (rural, suburban or urban) and the aesthetic, social, and safety aspects of the area to determine the vegetation selection, arrangement, and management.

Observed species selection criteria to improve aesthetics include seasonal foliage color, showy flowers, foliage texture, form, and branching habit. The layout and design should be appropriate for the setting as determined by adjacent land uses. A landscape analysis can help determine specific aesthetic requirements.

Where feasible, consider alternative water sources, such as tanks, ponds, wells, solar pumps, and ram pumps for livestock water supply needs.

Allelopathic impacts of plants should be considered.

PLANS AND SPECIFICATIONS

Specifications for applying this practice shall be prepared for each site. Specifications shall be recorded using approved specifications sheets, job sheets, narrative statements in the conservation plan, or other acceptable documentation. ***Requirements for operation and maintenance of the practice shall be incorporated into site specifications.***

The following will be in the conservation plan narrative or in the contract narrative (as appropriate):

Purpose of riparian forest buffer

Size of riparian forest buffer

- width of floodplain (ft.)
- width of riparian buffer (ft.)
- length of riparian buffer (ft.)
- acres of riparian buffer (ac.)

Field location / Plan view

Site conditions prior to establishment of the riparian forest buffer

Soil amendments (if needed)

Site preparation methods (if used)

Plant species

Plant Guides

Plant Sheets

Stock size

Spacing

Planting method(s)

Planting dates

Competition suppression

Protection methods

Cultural practices (i.e. pruning, forest stand improvement, etc.) (if any)

Alternative water source location

Fencing

Erosion control - blanket/mulch

Operation and Maintenance Plan

Replacement strategies

OPERATION AND MAINTENANCE

The following actions shall be carried out to insure that this practice functions as intended throughout its expected life.

These actions include normal repetitive activities in the application and use of the practice (operation), and repair and upkeep of the practice (maintenance)

The riparian forest buffer will be inspected periodically and protected to maintain the intended purpose from adverse impacts such as excessive vehicular and pedestrian traffic, pest infestations, pesticide use on adjacent lands, livestock damage, and fire.

The buffer should be inspected at least annually and after heavy storm events.

Check for areas where water is concentrated. Disperse concentrated flow by appropriate measures, including placement and repositioning debris.

Vehicular traffic or excessive animal traffic, and the removal or disturbance of vegetation and leaf litter must be avoided.

Where practical, management activities will be performed outside the primary nesting season (March 15 - July 15). An exception may be for mowing or cultivation to control vegetative competition.

Replacement of dead trees or shrubs and control of undesirable vegetative competition will be continued until the buffer is, or will progress to, a fully functional condition.

As applicable, control of concentrated flow erosion and sediment deposition shall be controlled by an adjacent filter strip.

Any removals of tree and shrub products shall be conducted in a manner that maintains the intended purpose.

Any use of fertilizers, pesticides and other chemicals to assure buffer function shall not compromise the intended purpose.

For the purpose of moderating water temperatures and providing detritus and large woody debris, riparian forest buffer management must maintain a minimum of 50 percent canopy cover.

For the purposes of reducing excess pollutants in surface runoff and shallow groundwater or providing habitat and corridors for wildlife, manage the canopy to maintain maximum vigor of the overstory and understory species.

Additional operation and maintenance requirements shall be developed on a site-specific basis to assure performance of the practice as intended.

REFERENCES

Castelle, A.J., A.W. Johnson, and C. Conolly. 1994. Wetland and Stream Buffer Size Requirements - A Review. Journal of Environmental Quality - Vol. 23 No. 4. Madison, WI.

Chesapeake Bay Program - Nutrient Subcommittee. 1995. Water Quality Functions of Riparian Forest Buffer Systems in the Chesapeake Bay Watershed. Prepared by: R. Lowrance, L.S. Altier, R.S. Schnabel, P.M. Groffman, J.M. Denver, D.L. Correll, J.W. Gilliam, J.L. Robinson, R.B. Brinsfield, K.W. Staver, W. Lucas, and A.H. Todd. Annapolis, MD.

Cornell Cooperative Extension Service. Wildlife and Timber from Private Lands: A Landowner's Guide to Planning. Info. Bull. 193. Prepared by: D.J. Decker, J.W. Kelley, T.W. Seamans, and R.R. Roth. Ithaca, NY. Gilliam, J.W. 1994. Riparian Wetlands and Water Quality. Journal of Environmental Quality - Volume 23: 896-900.

Schultz R.C., J.P. Colletti, T.M. Isenhardt, W.W. Simpkins, C.W. Mize, and M.L. Thompson. 1995. Design and Placement of a Multi-species Riparian Buffer Strip. Agroforestry Systems 29:201-225.

U.S. Department of Agriculture, Forest Service, Natural Resources Conservation Service, National Agroforestry Center. 1997. Riparian Buffer for Agricultural Land. Agroforestry Note #3. Prepared by: M. Doskey, D. Schultz, T. Isenhardt. Lincoln, NE.

U.S. Department of Agriculture, Forest Service, Natural Resources Conservation Service, National Agroforestry Center. 1997.

How to Design a Riparian Buffer for Agricultural Land. Agroforestry Note #4. Prepared by: M. Doskey, D. Schultz, T. Isenhardt. Lincoln, NE.

U.S. Department of Agriculture, Forest Service, Natural Resources Conservation Service, National Agroforestry Center. 1997. A Riparian Buffer Design for Cropland. Agroforestry Note #5. Prepared by: M. Doskey, D. Schultz, T. Isenhardt. Lincoln, NE.

U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry. 1993. Crop Tree Management in Riparian Zones. Forest Resources Management.

U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry. 1991. Riparian Forest Buffers - Function and Design for Protection and Enhancement of Water Resources. NA-PR-07-91. Prepared by David J. Welsch, Radnor, PA.

U.S. Department of Agriculture, Forest Service, Southern Region. 1992. Stream Habitat Improvement Handbook. Tech. Pub. R8-TP 16. Prepared by Monte E. Sheehorn, Atlanta, GA.

Welch, David E., D.L. Smart, J.N. Boyer, P. Minkin, H.C. Smith, T.L. McCandless. 1995. Forested Wetlands - Functions, Benefits, and the use of Best Management Practices. NA-PR-01-95.

West Virginia Division of Forestry. 1996. Best Management Practices for Controlling Soil Erosion and Sedimentation from Logging Operations in West Virginia. WVDOP-TR-96-3. August 2002, Charleston, WV.

**NATURAL RESOURCES CONSERVATION SERVICE
WEST VIRGINIA CONSERVATION PRACTICE STANDARD**

**RIPARIAN FOREST BUFFER
(Acre)**

CODE 391

Sample Practice Narratives

391-A- Riparian Forest Buffer - A buffer meeting at least the minimum width (at least 35 feet) requirement will be established and maintained along the stream corridor and/or water body. The landowner will devote this area to trees and/or shrubs by allowing natural regeneration to take place. This area will not be harvested, mowed or grazed. Control noxious weeds, such as thistle, multiflora rose, autumn olive or tartarian honeysuckle; by spot mowing or chemical weed control on areas where these species will interfere with natural succession. Do not mow or chemically treat seedling or sapling trees to ensure natural succession will result in establishment of the desired community. Mowing or chemical applications should not occur during the primary ground-nesting season (March 15 - July 15). Recommendations prescribed by WVU Extension Service, WV Division of Forestry personnel or as outlined in the Penn State Agronomy Guide will be followed. All applications of pesticides and/or herbicides will be consistent with state and federal guidelines and label precautions. Consult the attached specifications and/or job sheets for details and components necessary for establishment and maintenance of this practice.

Optional - In addition to the minimum combined widths of Zones 1 and 2 (35 feet) Zone 3 may also be established. This zone may be mowed and fertilized (not harvested) to maintain a 20-foot wide perennial sod/herbaceous cover along the inside of any planned or existing fence.

391-B- Riparian Forest Buffer - A buffer meeting at least the minimum width (at least 35 feet) requirement will be established and maintained along the stream corridor and/or water body. The landowner will devote this area to trees and/or shrubs by planting. This area will not be harvested, mowed or grazed. Control woody noxious weeds, such as multiflora rose, autumn olive or tartarian honeysuckle; by spot mowing or chemical weed control on areas where these species will interfere with planting or seedlings. Do not mow or chemically treat seedling or sapling trees to ensure natural succession will result in establishment of the desired community. Mowing or chemical applications should not occur during the primary ground-nesting season (March 15 - July 15). Recommendations prescribed by WVU Extension Service, WV Division of Forestry personnel or as outlined in the Penn State Agronomy Guide will be followed. All applications of pesticides and/or herbicides will be consistent with state and federal guidelines and label precautions. Consult the attached specifications and/or job sheets for details and components necessary for establishment and maintenance of this practice.

Optional - In addition to the minimum combined widths of Zones 1 and 2 (35 feet) Zone 3 may also be established. This zone may be mowed and fertilized (not harvested) to maintain a 20-foot wide perennial sod/herbaceous cover along the inside of any planned or existing fence.